arXiv:1301.7226v2 [cond-mat.quant-gas] 5 Mar 2013

Triplon Supersolid of Cold Atoms in a Ladder-Shaped Optical Lattice

Qi-Hui Chen[1,2], Peng Li[2*], Han-Ting Wang[1] and Wu-Ming Liu[1†]
[1] *National Laboratory for Condensed Matter Physics, Institute of Physics,*
Chinese Academy of Sciences, Beijing 100190, China and
[2] *The Key Laboratory of High Energy Density Physics and Technology*
(Ministry of Education), Sichuan University, Chengdu 610064, China

(Dated: March 6, 2013)

We investigate the ground state phase diagram of the cold atoms in a ladder-shaped optical lattice by using stochastic series expansion quantum Monte Carlo method. We demonstrate how triplon supersolid could emerge, in which the triplon superfluid can coexist with the long-range triplon solid. In the triplon superfluid the quasi-condensation appears in the form of power-law decaying phase correlations. We suggest that a quasi-supersolid phase could be charted in the phase diagram, in which both superfluid and solid orders are quasi-long range ones. We also discuss how this phenomenon can be realized and detected in real experiment.

PACS numbers: 03.75.Lm, 03.75.Nt, 37.10.Jk, 67.80.kb

Introduction.—With the advantage of cleanness and controllability, the cold atoms system in optical lattice has been a powerful platform to simulate a variety of quantum system. Many works in experimental and theoretical aspects have been made in the past few decades [1–6]. Various types of optical lattices with interesting geometric structure, such as triangular, kagome and honeycomb [7–9], could be realized by adjusting the potential formed by lasers. Further more, the interactions between atoms can also be tuned by Feshbach resonance [10]. With the aid of ultracold gas in optical lattice, the quantum phase transition from superfluid to Mott insulator is also observed [1]. Beyond that, we ask whether some fascinating quantum states, such as supersolid (SS) which is not observed in real materials so far, could be implemented by using cold atoms in some new kinds of optical lattices, such as a ladder-shaped optical lattice.

Featured by the coexistence of both diagonal and off-diagonal long-range order, SS is an intriguing exotic phase of matter, of which research works date back to fifty year ago. Aroused mainly by the tentative experimental evidence for the presence of superfluid (SF) in solid ^{4}He [11], although it is confirmed that all of those early results were misunderstood recently [12], an upsurge in quest for supersolid has been lasting for nearly a decade [13–21]. In another route, lattice SS have been paid much attention, especially after the advances in optical lattice technology [10, 22]. Various schemes for lattice systems from one to three dimensions have been designed [23–30]. Recently, the cold atoms in isolated double-well have been realized in experiment [2–5]. Could we go beyond those experiments to design a new system and realize a novel SS phase in optical lattices?

In this Letter, we investigate SS phase by extending the isolated double-well [2–5] to a spin dependent ladder-shaped optical lattice, as illustrated in Fig. 1. It's can be viewed as double-well lattice that are coupled along y direction. Comparing with real compounds, the advantage of this system are the cleanness and highly controllability of the interactions which allow us extend previous studies of ladder system [31–33] to the parameters region where the triplon SS can emerge. This novel phase can be observed in ladder-shaped optical lattice by varying the lattice depth, the zeeman field and the interaction strength via the Feshbach resonance. When the interactions between cold atoms is larger than its kinetic enough i.e., $U \gg t$, the system enter Mott insulator region. The dynamics of atoms just involves spin freedom of degree. By adjusting the zeeman field and the hopping of cold atoms with different spin, the triplon SF is induced by the power-law decaying phase correlations, while the triplon solid emerges as the result of triplon interactions. In a proper parameters region, triplon SF and solid appear simultaneously to form the triplon SS.

The cold atoms in a ladder-shaped optical lattice.—A spin dependent ladder-shaped optical potential is constructed as Fig. 1. We assume that only two of the internal ground states participate in the dynamics, which we call spin up and down, $|\uparrow\rangle$ and $|\downarrow\rangle$. By second-order perturbation theory for coupling of the ground state levels to the excited atomic states, the potential can be obtained as, $U_\sigma = V_{x1,\sigma}\cos^2(k_x x) + V_{x2,\sigma}\cos^2(2k_x x) + V_{y,\sigma}\cos^2(k_y y)$, where $\sigma = \uparrow, \downarrow$ represents the two states of cold atoms and $V_{x1,\sigma}$ and $V_{x2,\sigma}$ are the barrier heights of the two standing waves along the x direction, $V_{y,\sigma}$ is the one along the y direction. In experiment a harmonic potential with frequency ω_z should be added in z direction which restricts the moving of atoms along it. To simulate a decoupled two-leg ladder potential, $4V_{x2,\sigma} > V_{x1,\sigma}$ is required and the ratio of potential heights between intra-ladder to inter-ladder along x direction should satisfies $\Delta V = (4V_{x2,\sigma} - V_{x1,\sigma})^2/(4V_{x2,\sigma} + V_{x1,\sigma})^2 \ll 1$. The widths of the two barriers along rung direction and rail direction are $a = \alpha/k_x$ and $b = \pi/k_y$, respectively, with $\alpha = \arccos(V_{x1,\sigma}/4V_{x2,\sigma})$. With the considerations, then by using harmonic approximation in each bottom of trap,



FIG. 1: (Color online) A ladder-shaped optical potential is formed by three standing wave lasers indicated as a pair of counter yellow arrows V_{x1}, V_{x2}, V_y. Two V_{x1}, V_{x2} of them along x direction and another one V_y along y direction, where V_{x2} (short yellow arrows) has twice the period of V_{x1} (long yellow arrows). The cold atoms with two internal states denoted by spin up $|\uparrow\rangle$ (red up arrows) and spin down $|\downarrow\rangle$ (blue down arrows) are trapped by this potential. An antiferromagnetic (AF) configuration with staggered up-and-down moments on the ladder is showed. The white lines in the potential contour lines represent the ladder. The effective strength of interactions is $J_\perp$ along x direction, $J^x_\parallel, J^z_\parallel$ along y direction. h is zeeman field.



FIG. 2: (Color online) Ground state phase diagram with $J^x_\parallel = 0.35$. The doted lines (Gray) are results from the mean field theory. The scatter lines (Black) are QMC results. There are four orders which are antiferromagnetic (AF), full polarized (FP), and triplon solid (TS) orders as shown in the insets. The triplon superfluid (SF) phase occupies a considerable area. The triplon supersolid (SS) phase is sandwiched between the AF and TS phase. The quasi-SS (q-SS) phase, which exhibits algebraically decaying diagonal TS order, is also charted.

this system can be described by Hubbard model as,

$$H = \sum_{i,j,\sigma}(-t_{\mu,\sigma}c^\dagger_{i,\sigma}c_{j,\sigma} + h.c) + U_{\uparrow\downarrow}\sum_i n_{i,\uparrow}n_{i,\downarrow} + \sum_{i,\sigma}U_\sigma n_{i,\sigma}(n_{i,\sigma}-1) - \tilde{h}\sum_i(n_{i,\uparrow}-n_{i,\downarrow}), \tag{1}$$

where a zeeman term $\tilde{h}$ is also introduced. $c^\dagger_{i,\sigma}(c_{i,\sigma})$ is the creation (annihilation) operator of boson or fermion in site i with spin σ. The coefficients $t_{\mu,\sigma}$, with $\mu = \perp, \parallel$, represent intraladder hopping along rung, rail direction, respectively. $U_{\uparrow\downarrow}, U_\sigma$ are the on-site interaction between atoms. We can work out those interaction parameters as a function of $V_{x1,\sigma}, V_{x2,\sigma}, V_{y,\sigma}, a_s, E_{xr}, E_{yr}$, where a_s is the scattering lengths and $E_{x,yr} = \hbar^2k^2_{x,y}/2m$ are the atomic recoil energy along x, y direction respectively.

Considering $t_\mu \ll U_{\uparrow\downarrow}, U_\sigma$ and 1/2 filling, by a second order perturbation theory, this system will be described by an effective Hamiltonian with spin freedom of degree only, $\tilde{H} = \sum_{i,j}[J^z_\mu S^z_i S^z_j \pm J^x_\mu(S^x_i S^x_j + S^y_i S^y_j)] - h\sum_i S^z_i$, where $\mathbf{S} = \frac{1}{2}c^\dagger\vec{\sigma}c$, $\vec{\sigma}$ is pauli matrix and $c^\dagger = (c^\dagger_\uparrow, c^\dagger_\downarrow)$. The upper sign before J^x_μ is for fermionic atoms and down sign for bosonic one. The interaction coefficients for bosons can be figured out as, $J^z_\mu = \frac{2(t^2_{\mu\uparrow}+t^2_{\mu\downarrow})}{U_{\uparrow\downarrow}} - \frac{t^2_{\mu\uparrow}}{U_\uparrow} - \frac{t^2_{\mu\downarrow}}{U_\downarrow}$, $J^x_\mu = \frac{4t_{\mu\uparrow}t_{\mu\downarrow}}{U_{\uparrow\downarrow}}$, $h = 2\tilde{h} + \frac{6t^2_{\mu\uparrow}}{U_\uparrow} - \frac{6t^2_{\mu\downarrow}}{U_\downarrow}$. For fermion, we need to omit the last two term in J^z_μ and h.

Since the lattice is bipartite, without loss of generality, we focus on the region with positive J^x_μ of the spin XXZ model $\tilde{H}$ and hereafter we take $J^x_\perp = J^z_\perp = J_\perp = 1$ as energy unit. The QMC simulation [34] is performed on a ladder with length L and temperature $T = 6/L$. A mean-field (MF) theory is also employed for comparison. The ground-state phase diagram is illustrated in Fig. 2.

The triplon superfluidity.—Let's start from the isotropic case, i.e., $J^x_\parallel = J^z_\parallel = J_\parallel$, which has been extensively studied in the context of real materials [35]. The four eigenstates of $j\,th$ decoupled rung provide very helpful basis: the singlet state, $|s\rangle_j = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle_j - |\downarrow\uparrow\rangle_j)$, and three triplet states, $|t_+\rangle_j = |\uparrow\uparrow\rangle_j$, $|t_0\rangle_j = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle_j + |\downarrow\uparrow\rangle_j)$, $|t_-\rangle_j = |\downarrow\downarrow\rangle_j$. If no (or weak) magnetic field is present, the ladder is in a gapped SD phase, whose ground state is roughly a chain of singlets. While the system will be in a FP phase in the strong magnetic field limit. The SD and FP phases can be readily expressed by MF wave functions, $|\psi_{\mathrm{SD}}\rangle = \prod_j |s\rangle_j$ and $|\psi_{\mathrm{FP}}\rangle = \prod_j |t_+\rangle_j$, respectively. If moderate field is applied, a gapless triplon SF phase emerges, which is induced by the quasi-condensation of tripon when the energy level of $|t_+\rangle$ is down to the vicinity of state $|s\rangle$. And the upper $|t_0\rangle$ and $|t_-\rangle$ bands could be integrated out from the renormalization point of view. One can introduce triplon bosonic operators $\widetilde{b}^\dagger_j = \frac{1}{\sqrt{2}}(S^+_{j,2} - S^+_{j,1})$ that fulfill $\widetilde{b}^\dagger_j|s\rangle_j = |t_+\rangle_j$. Then the system could be mapped to a 1D hardcore bose-Hubbard model. It can be exactly solved by Bethe Ansatz or approximately by Bosonization method, gives the Luttinger liquid (LL) nature [36]. In this regime, the state $|t_+\rangle$ undergoes a quasi-condensation characterized by a power-law decaying phase correlations $\langle\widetilde{b}^\dagger_j\widetilde{b}_{j+r}\rangle \sim r^{-1/2K}\cos\pi r$, where K is the LL parameter. Although true condensation is absent in 1D systems, the triplon superfluidity is still possible due to the power-law decaying phase cor-



FIG. 3: (Color online) (a) The desity peak in momentum space $\langle\widetilde{n}_{\pi}\rangle$, (b) The triplon structure factor $S\left(\pi\right)$ for different system sizes, (c) The extracted index α and β versus h for the isotropic case $J_{\parallel}^{x}=J_{\parallel}^{z}=0.35$. (d) The staggered magnetization m_{s} and (e) its corresponding staggered magnetic susceptibility χ_{s} for $J_{\parallel}^{x}=0.35$ and $h=0.5$.

relations. The (sublinearly) diverging peak [29] of the momentum distribution $\langle\widetilde{n}_{k}\rangle=\frac{1}{L}\sum_{j,r}e^{ikr}\langle\widetilde{b}_{j}^{\dagger}\widetilde{b}_{j+r}\rangle$ occurs at $k=\pi$ and the quasi-condensation index α can be extracted from the formula $\langle\widetilde{n}_{\pi}\rangle\propto L^{\alpha}(\alpha<1)$. The calculation of $\langle\widetilde{n}_{\pi}\rangle$ is carried out in SSE frame with algorithm of measuring Green's function [37].

In Fig. 3 (a-c), we show the density peak $\langle\widetilde{n}_{\pi}\rangle$, the structure factor $S(\pi)$ and the index α versus h. In there, the transition points are determined by the occurring of meaningful nonzero α, if the fitted $\alpha\leq 0$, that means there is other order instead of the triplon superfluid, thus we denote it as zero. To check the validity of the use of quasi-condensation index α, we work out the transition points at the isotropic case. For $J_{\parallel}^{x}=J_{\parallel}^{z}=0.35$, SSE simulation for the original ladder system gives the transition points from SD to SF, $h_{c1}=0.65\pm 0.03$), and from SF to FP, $h_{c2}=0.88\pm 0.02$) (see Fig. 3 (c)). The exact solutions of it are $h_{c1}=J_{\perp}-J_{\parallel}=0.65$ and $h_{c2}=J_{\perp}+2J_{\parallel}=0.85$, in good agreement with QMC.

The triplon supersolid.—When we raise the anisotropy along the rail [39], say $J_{\parallel}^{x}\neq J_{\parallel}^{z}$, the SF spreads to a considerable area, which can be charted by the quasi-condensation index α. If the anisotropy is large enough, two types of Ising orders are energetically favored. The first one is the traditional Ising order, i.e., AF arrangement. It is the usual Ising order and can be represented by $|\psi_{\rm AF}\rangle=\prod_{j}(u|s\rangle_{j}+(-1)^{j}w|t_{0}\rangle_{j})$ with u and w being the variational parameters to minimize the MF energy. The order parameter is the staggered magnetization $m_{s}=\frac{1}{N}\sum_{j}\langle(-1)^{j}(S_{j,1}^{z}-S_{j,2}^{z})\rangle$. The transition from SD to AF is of second order, which is well signified by the divergent peak of the staggered susceptibility $\chi_{s}=\frac{\partial m_{s}}{\partial h}$ (see Fig. 3(d) and (e)).



FIG. 4: (Color online) (a) The density peak $\langle\widetilde{n}_{\pi}\rangle$ and (b) the structure factor $S(\pi)$ versus L in log-log scale. The parameters are: SF ($h=0.96,\alpha=0.528\pm 0.001$), q-SS ($h=1.2,\alpha=0.513\pm 0.004,\beta=0.177\pm 0.002$) in isotropic case $J_{\parallel}^{z}=0.35$, TS ($h=4.2,\beta=1.433\pm 0.009$), SS ($h=4.06,\alpha=0.373\pm 0.008,\beta=1.082\pm 0.003$) in anisotropic case $J_{\parallel}^{z}=4.0$. The dotted lines have slope value 1 which facilitates to discern the occurence of true TS order.

The second type of solid order is the alternative arrangement of singlet state $|s\rangle$ and triplet state $|t_{+}\rangle$ on the rung along the rail of the ladder. It is the TS order represented by $|\psi_{\rm TS}\rangle=\prod_{j}(u_{A}|s\rangle_{2j}+(-1)^{j}v_{A}|t_{+}\rangle_{2j})\otimes(u_{B}|s\rangle_{2j+1}+(-1)^{j}v_{B}|t_{+}\rangle_{2j+1})$ with variational parameters $u_{A}\neq u_{B}$ and $v_{A}\neq v_{B}$. The TS order breaks the translational symmetry of the ladder along the rail. To detect the TS order, we use the triplon structure factor $S(\pi)=\frac{1}{L}\sum_{i,j}(-1)^{(i+j)}\langle\widetilde{n}_{i,t_{+}}\widetilde{n}_{j,t_{+}}\rangle$. True TS order means a finite diagonal order parameter squared, $\langle m_{\rm TS}\rangle^{2}=\frac{1}{L}S(\pi)$. The true TS order is a consequence of the interplay between the anisotropy and the strong field. While in the SF regime, there may exist a weak quasi-triplon solid order that arises from the power-law decaying correlators [36]. It could be further strengthened by the anisotropy. So we define the diagonal order index β that shall be extracted by the formula $S(\pi)\propto L^{\beta}$ and set up the conditions for judging diagonal TS orders: if $\beta\gtrsim 1$, there exists a true TS order; if $\beta<1$, a quasi-TS order; while if no suitable β, no any TS order. Meanwhile, if superfluidity (i.e., persistent LL) also survives under the anisotropy, then one would possibly get true SS order as well as q-SS order.

We demonstrate the density peak $\langle\widetilde{n}_{\pi}\rangle$ and the structure factor $S(\pi)$ versus lattice size in log-log scale in Fig. 4. From the slope of the data lines, one can figure out the indices α and β. It is definite to see the coexistence of TS order and quasi-condensation in the SS phase. We also noted that the q-SS phase can occur at the isotropic case, as is shown in Fig. 3 (c). We show the density peak $\langle\widetilde{n}_{\pi}\rangle$, the structure factor peak $S(\pi)$ in different lattice sizes, the indices α and β versus h (see Fig. 5). We see the SS phase is a intermediate stage of the transition from TS to AF phase. In TS phase, the densities n_{s} and n_{t+}



FIG. 5: (Color online) (a) The density peak $\langle\tilde{n}_\pi\rangle$ and (b) the structure factor $S(\pi)$ versus external field h for different system size L. (c) The index α and β fitted from $\tilde{n}(\pi)$ and $S(\pi)$ where the model parameters are $J_x = 0.35$, $J_z = 4.0$.



FIG. 6: (Color online) Histograms of m_z of a small area (a-d) of size $l = 8$ (red), $l = 12$ (green) and the whole lattice (e-h) of size $L = 88$ (Blue). The parameters are $J_z = 2.0$ and $h = 2.03, 2.17, 2.23, 2.53$ for SS, TS, q-SS, and SF, respectively.

prevail. Before entering into the AF phase, we find the system gains a considerable small fraction of $n_{t0} \sim 0.13$, which plays the role of impurities in the SS phase [26]. This may be the reason why SS can not emerge in general 1D hard-core boson model or XXZ model but in this system. It is worthwhile to note that the simple mean-field description for the superfluidity in 1D is not quite right. One can combine all the components in a wave function as $|\psi\rangle = \prod_j (u|s\rangle_j + (-1)^j v|t_+\rangle_j + (-1)^j w|t_0\rangle_j)$. But this assumption would lead to a true condensation and, thus, totally failed to capture the quasi-condensation nature. So the situation here is quite similar to the 1D spin 1 chain with uniaxial exchange and single-ion anisotropies, where the LL behavior also plays an important role in providing the 1D superfluidity [28].

It is important to preclude the possibility of phase separation. This can be done by measuring m_z of the small area of size (say $l = 8(12)$) throughout the whole lattice ($L = 88$). If no phase separation occurs, we should observe only a single peak in the histograms of m_z [26]. This is indeed the case as shown in Fig. 6.

Experimental proposal.—The system discussed in this paper may be realized in a Mott insulating phase of cold bosonic or fermionic atoms in a ladder-shaped optical lattice if the atoms possess two internal states that play the role of the spin (such as ^{87}Rb in $|\uparrow\rangle \equiv |F = 1, m_F = 1\rangle, |\downarrow\rangle \equiv |F = 1, m_F = -1\rangle$). Let's estimate the typical energy scales. For Rb atoms with a lattice constant $\pi/k_x \simeq 2\pi/k_y \sim 426$ nm and about 10^5 atoms in a Bose-Einstein condensate, we can chose $t^2/(\hbar U) \sim 0.1$ kHz (corresponding to a time scale of 10 ms) with a conservative choice of $U \sim 2$ kHz and $(t/U)^2 \sim 1/20$. These energy scales are clearly compatible with current experiments [1] and make the system in a Mott insulating area.

In experiment, the density of condensates in momentum space $\langle\tilde{n}_\pi\rangle$ can be measured by noise correlations which can be linked to spin-spin correlations [41–43]. We can use Bragg scattering of light, which gives rise to the spin structure factor, to detect $S(\pi)$ [44]. An alternative technique for imaging spin states in optical lattices has been put forward [45]. Thus, the SS discussed in this paper can be detected in experiment.

Conclusions.—In summary, we propose a scheme to realize triplon SS of cold atoms in a ladder-shaped optical lattice. There is a q-SS phase emerging as a precursor of the true SS phase and TS phases. The most importance is that this triplon SS can be detected in recent experimental technology. These results are beyond both previous experimental researches in isolated double-well and previous studies of ladder system. It can be used as a guide for future experimental studies of supersolid and quasi-supersolid.

This work was supported by the NKBRSFC under grants Nos. 2011CB921502, 2012CB821305, 2009CB930701, 2010CB922904, NSFC under grants Nos. 10934010, 11074177, 11228409, 61227902, NSFC-RGC under grants Nos. 11061160490 and 1386-N-HKU748/10, SRF for ROCS SEM (20111139-10-2).

---

[1] M. Greiner et al., Nature (London) **415**, 39 (2002).
[2] S. Trotzky et al., Science **319**, 295 (2008).
[3] S. Fölling et al., Nature (London) **448**, 1029 (2007).
[4] Y. A. Chen, S. Nascimbéne, M. Aidelsburger, M. Atala, S. Trotzky, and I. Bloch, Phys. Rev. Lett. **107**, 210405 (2011).
[5] J. Sebby-Strabley, M. Anderlini, P. S. Jessen, and J. V. Porto, Phys. Rev. A **73**, 033605 (2006).
[6] L. M. Duan, E. Demler and M. D. Lukin, Phys. Rev. Lett. **91**, 090402 (2003).

[7] L. Santos et al., Phys. Rev. Lett. **93**, 030601 (2004).

[8] P. Soltan-Panachi et al., Nature Phy. **7**, 434 (2011).

[9] Y. H. Chen, W. Wu, H. S. Tao and W. M. Liu, Phys. Rev. A **82**, 043625 (2010).

[10] I. Bloch, J. Dalibard and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).

[11] E. Kim and M. H. W. Chan, Nature (London) **427**, 225 (2004); Science **305**, 1941 (2004).

[12] D. Y. Kim and M. H. W. Chan, Phys. Rev. Lett. **109**, 155301 (2012).

[13] A. Koga, T. Higashiyama, K. Inaba, S. Suga and N. Kawakami, Phys. Rev. A **79**, 013607 (2009).

[14] A. Bühler and H. P. Buchler, Phys. Rev. A **84**, 023607 (2011).

[15] D. Yamamoto, I. Danshita and Carlos A. R. Sá de Melo, Phys. Rev. A **85**, 021601(R) (2012).

[16] I. Danshita and Carlos A. R. Sá de Melo, Phys. Rev. Lett **103**, 225301 (2009).

[17] K. P. Schmidt, J. Dorier and A. M. Lauchli, Phys. Rev. Lett. **101**, 150405 (2008).

[18] J. W. Ye, Phys. Rev. Lett. **97**, 125302 (2006).

[19] Y. H. Chan, Y. J. Han and L.M. Duan, Phys. Rev. A **82**, 053607 (2010).

[20] X. P. Li, W. V. Liu and C. W. Lin, Phys. Rev. A **83**, 021602(R) (2011).

[21] P. P. Orth, D. L. Bergman and K. Le Hur, Phys. Rev. A **80**, 023624 (2009).

[22] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner and P. Zoller, Phys. Rev. Lett. **81**, 3108 (1998).

[23] S. Wessel and M. Troyer, Phys. Rev. Lett. **95**, 127205 (2005).

[24] D. Heidarian and K. Damle Phys. Rev. Lett. **95**, 127206 (2005).

[25] R. G. Melko, A. Paramekanti, A. A. Burkov, A. Vishwanath, D. N. Sheng and L. Balents, Phys. Rev. Lett. **95**, 127207 (2005).

[26] K. K. Ng and T. K. Lee, Phys. Rev. Lett. **97**, 127204 (2006).

[27] S. Yi, T. Li, and C. P. Sun, Phys. Rev. Lett. **98**, 260405 (2007).

[28] P. Sengupta and C. D. Batista, Phys. Rev. Lett. **98**, 227201 (2007); *ibid*, **99**, 217205 (2007).

[29] T. Keilmann, I. Cirac, and T. Roscilde, Phys. Rev. Lett. **102**, 255304 (2009).

[30] L. Mathey, I. Danshita and C. W. Clark, Phys. Rev. A **79**, 011602(R) (2009).

[31] Y. Fujihara, A. Koga and N. Kawakami, Phys. Rev. A **79**, 013610 (2009).

[32] F. Crepin, R. Citro and P. Simon, Phys. Rev. A **82**, 013613 (2010).

[33] M. Okumura, S. Yamada, M. Machida and H. Aoki, Phys. Rev. A **83**, 031606(R) (2011).

[34] O. F. Syljuasen and A. W. Sandvik, Phys. Rev. E **66**, 046701 (2002).

[35] E. Dagotto and T. M. Rice, Science **271**, 5249 (1996).

[36] T. Giamarchi, *Quantum Physics in One Dimension* (Clarendon, Oxford 2003).

[37] A. Dorneich and M. Troyer, Phys. Rev. E **64**, 066701 (2001).

[38] N. Laflorencie and F. Mila, Phys. Rev. Lett. **99**, 027202 (2007).

[39] One can also consider the anisotropy on the rung. But to produce the triplon SS phase we concern, this anisotropy is unnecessary.

[40] In Ref. [29], this phase is attributed to “s-Tonks” phase. Here we adopt the term “quasi-supersolid”.

[41] E. Altman, E. Demler, and M. D. Lukin, Phys. Rev. A **70**, 013603 (2004).

[42] S. Fölling et al., Nature (London) **434**, 481 (2005).

[43] V. W. Scarola, E. Demler and S. DasSarma, Phys. Rev. A **73**, 051601(R) (2006).

[44] T. A. Corcovilos, S. K. Baur, J. M. Hitchcock, E. J. Mueller and R. G. Hulet, Phys. Rev. A **81**, 013415 (2010).

[45] J. S. Douglas and K. Burnett, Phys. Rev. A **82**, 033434 (2010).